

# Decoherence in quantum field theory


Julien Serreau

APC, Univ. Paris Diderot

[ A. Giraud, J.S. : submitted to PRL ]

- Motivations and set-up
- Nonequilibrium QFT: methods
- Results
- Conclusions and outlook

# MOTIVATIONS



# The standard picture

Decoherence : dynamical diagonalization of the density matrix (in a given basis)

ex:

$$\rho(t=0) = |\psi\rangle\langle\psi|$$

pure state

$$\rho(t) = \sum_n p_n |n\rangle\langle n|$$

statistical mixture




Loss of purity / coherence results from the interaction with an "environment" or from some kind of coarse graining

NON-UNITARY  
EVOLUTION

# Decoherence for a closed system

- Reconstructing the actual state of the system requires one to measure all independent correlation functions.
- For systems with large (infinite) number of d.o.f., this is practically impossible.

➔ The inability to reconstruct the actual state of the system due to restricted data may result in apparent decoherence.



Apparent "trajectory" in the subspace of measured correlation functions

Information spreads in the space of correlation functions.

[ e.g. Balian ('00); Campo, Parentani ('05) ... ]

# Theoretical description

Decoherence: a nonequilibrium quantum process.

## ■ Analytical descriptions

- exactly solvable (simple) models
- neglect the dynamics of the environment (e.g. assume a thermal bath)
- neglect back-reaction of the system
- weak coupling techniques

## ■ Numerical work

- solve open quantum dynamics obtained by integrating out the environment d.o.f.



## Real-time quantum dynamics

# Nonequilibrium QFT : methods

⚠ Standard approximation schemes in QFT fail out of equilibrium.

● SECULARITY :  $e^{-\gamma t} \approx 1 - \gamma t + \dots$

⇒ Need for (infinite) resummations

● UNIVERSALITY : late-time thermalization (effective loss of memory)

⇒ Need for non-linear approximation schemes

Two-particle-irreducible (2PI) techniques

[e.g. Serreau @ SEWM08 hep-ph/0410330]

$$\Gamma_{2PI}[\phi, G] = S[\phi] + \frac{i}{2} \text{Tr} \text{Ln} G^{-1} + \frac{i}{2} \text{Tr} G_0^{-1} G + \Gamma_2[\phi, G]$$

$$\frac{\delta \Gamma_{2PI}}{\delta G} \Big|_{\bar{G}} = 0 \iff \bar{G}^{-1} = G_0^{-1} - \Sigma[\bar{G}]$$

Self-adjusting propagator

# 2PI techniques : State of art

➔ Solve the secularity and universality problems of nonequilibrium QFT. [Berges ('01), Cox ('01)]

## ■ Applications

- Thermalization from first principles [Berges, Borsanyi, J.S. ('03); Cooper, Dawson, Fluhica ('05)]
- Nonperturbative  $1/N$  expansion [Aarts, Ahrensmeier, Baier, Berges, J.S. ('02)]
- Dynamics at nonperturbatively high densities e.g. preheating after inflation [Berges, J.S. ('03), Smit, Tranberg ('04)]
- Expanding geometries [Tranberg ('08)]

⋮

## ■ Formal developments

- 2PI renormalization theory [Van Hees, Knoll ('02); Blaizot, Iancu, Reinosca ('03), Berges, Borsanyi, Reinosca, J.S. ('05) ...]
- Gauge field theories [Reinosca, J.S. ('06)]

⋮

# Decoherence in QFT : The model

- $O(N)$  scalar field  $\phi_a, a = 1 \dots N$

$$S[\phi] = \int d^4x \left[ \frac{1}{2} \partial_\mu \phi_a \partial^\mu \phi_a - \frac{m^2}{2} \phi_a \phi_a - \frac{\lambda}{4!N} (\phi_a \phi_a)^2 \right]$$

- 2PI  $\mathcal{K}_N$ -expansion at NLO [Berges ('02)]  
[Aarts, Ahrensmeier, Baier, Berges, J.S. ('02)]

$$\Gamma_2[G] = \Gamma_2^{LO}[G] + \Gamma_2^{NLO}[G] + \dots$$

$$\text{Loop diagram} \sim \frac{\lambda}{N} N^2 \sim \lambda N$$

$$\sum_{P \geq 3} \text{Diagram} \sim \frac{\lambda^P}{N^P} N^P \sim N^0$$

Gaussian dynamics : no "decoherence"

Non-Gaussian

$$\Sigma[G] = 2i \frac{\delta \Gamma_2}{\delta G} = \underbrace{\text{Loop diagram}}_{LO \text{ (mass correction)}} + \underbrace{\text{Diagram} + \dots + \text{Diagram}}_{NLO \text{ (scattering, memory effects, ...)}}$$



# Equations of motion

$$G(x,y) = \langle T \varphi(x) \varphi(y) \rangle = F(x,y) - \frac{i}{2} \varepsilon(x^0 - y^0) \rho(x,y)$$

statistical ↗
spectral ↗

$$[\square + M^2(x)] F(x,y) = - \int_0^{x^0} dz \Sigma_p(x,z) F(z,y) + \int_0^{y^0} dz \Sigma_F(x,z) \rho(z,y)$$

$$[\square + M^2(x)] \rho(x,y) = - \int_{y^0}^{x^0} dz \Sigma_p(x,z) \rho(z,y)$$

(local) mass corrections

Nonlocal terms  
= 'memory' / scatterings

$$\Sigma_F(x,y) = \frac{\lambda}{3N} \left[ F(x,y) \mathbb{I}_F(x,y) - \frac{1}{4} \rho(x,y) \mathbb{I}_\rho(x,y) \right]$$

$$\Sigma_p(x,y) = \frac{\lambda}{3N} \left[ F(x,y) \mathbb{I}_\rho(x,y) + \rho(x,y) \mathbb{I}_F(x,y) \right]$$

$$\mathbb{I}_F(x,y) = \mathbb{T}_F(x,y) - \int_0^{x^0} dz \mathbb{I}_\rho(x,z) \mathbb{T}_F(z,y) + \int_0^{y^0} dz \mathbb{I}_F(x,z) \mathbb{T}_\rho(z,y)$$


$$\mathbb{I}_\rho(x,y) = \mathbb{T}_\rho(x,y) - \int_{y^0}^{x^0} dz \mathbb{I}_\rho(x,z) \mathbb{T}_\rho(z,y)$$

$$\mathbb{T}_F(x,y) = -\frac{\Delta}{6} \left[ F^2(x,y) - \frac{1}{4} \rho^2(x,y) \right]; \quad \mathbb{T}_\rho(x,y) = -\frac{\Delta}{3} F(x,y) \rho(x,y)$$

# The picture

Assume one only measures equal-time 2-point functions:  $F_p(t) = \langle \varphi_p^\dagger(t) \varphi_p(t) \rangle$ ;  $K_p(t) = \langle \pi_p^\dagger(t) \pi_p(t) \rangle$

$$R_p(t) = \frac{1}{2} \langle \pi_p^\dagger(t) \varphi_p(t) + \varphi_p^\dagger(t) \pi_p(t) \rangle$$



The least-biased (effective) density matrix compatible with the known data:

$$\rho_{\text{eff}}(t) = \prod_p \rho_p(t)$$

$$\rho_p(t) \propto \exp \left[ K_p(t) \left( F_p(t) \pi_p^\dagger \pi_p + K_p(t) \varphi_p^\dagger \varphi_p + R_p(t) (\varphi_p^\dagger \pi_p + \pi_p^\dagger \varphi_p) \right) \right]$$

$$K_p(t) = - \frac{\ln(1 + 1/n_p(t))}{2n_p(t) + 1}; \quad n_p(t) + \frac{1}{2} = \sqrt{F_p(t)K_p(t) - R_p^2(t)}$$

N.B.:

$$\text{tr} \rho_p^2(t) = \frac{1}{2n_p(t) + 1}$$

: a measure of purity

# The Gaussian density matrix

Parametrization:

$$0 \leq \phi < 2\pi$$

$$0 \leq \gamma < 1$$


$$EF = \left(n + \frac{1}{2}\right) \frac{1 - \gamma \cos \phi}{\sqrt{1 - \gamma^2}}$$

$$R = -\left(n + \frac{1}{2}\right) \gamma \frac{\sin \phi}{\sqrt{1 - \gamma^2}}$$

$$K/\varepsilon = \left(n + \frac{1}{2}\right) \frac{1 + \gamma \cos \phi}{\sqrt{1 - \gamma^2}}$$

$\varepsilon$  = energy scale: fixes the basis  
we take  $\varepsilon_p(t) = \sqrt{p^2 + M^2(t)}$

$P_p(t)$



$$\sqrt{\left(n + \frac{1}{2}\right) \frac{1 - \gamma}{1 + \gamma}}$$

# Observables

$$\text{tr } \rho_p^2(t) = \frac{1}{2n_p(t)+1} \leq 1$$

↑  
Pure state  
( $n=0$ )

PURITY  
(basis-indep.)

$$S_p(t) = -\text{tr } \rho_p(t) \ln \rho_p(t)$$

$$= (n_p(t)+1) \ln(n_p(t)+1) - n_p(t) \ln n_p(t)$$


ENTROPY = MISSING INFORMATION  
(basis-indep.)

$\gamma_p(t)$ : squeezing parameter

COHERENCE in two-modes coherent states basis [Campo, Parentani ('05)]

# Initial conditions


We assume a Gaussian initial state



N.B.: No environment, no (thermal) bath...

Loss of quantum purity/coherence is to be caused by vacuum quantum fluctuations !!

# Results




Apparent loss of purity  $\epsilon_p$  and thermalization

The pure state looks mixed and even (later) thermal to the observer who has restricted information.


# Results

## Decoherence



$k = 0^+$

decoherence rate



Loss of quantum purity/coherence induced by quantum fluctuations

# Classical scaling

$$F, R, K \sim \frac{1}{\sqrt{1-\gamma^2}} \gg_{\gamma \rightarrow 1} p \sim 1$$

Classical (statistical) fluctuations regime

$$S[\varphi] = \int d^4x \left( \frac{1}{2} (\partial\varphi)^2 - \frac{m^2}{2} \varphi^2 - \frac{\lambda}{4!N} \varphi^4 \right)$$

$$= \eta \int d^4x \left( \frac{1}{2} (\partial\varphi')^2 - \frac{m^2}{2} \varphi'^2 - \frac{\lambda\eta}{4!N} \varphi'^4 \right)$$

$$\varphi = \sqrt{\eta} \varphi'$$

effective coupling

In the classical (strong) field regime, the results should only depend on

$$\lambda_{\text{eff}} = \frac{\lambda}{\sqrt{1-\gamma_0^2}}$$

Loss of purity/coherence can be described by means of classical statistical field theory




# Conclusions ...

- effective loss of quantum purity / coherence due to restricted knowledge on the system.
- induced by pure quantum fluctuations.
- can be described by classical statistical (stochastic) field theory.

# ... and Outlook

- ➔ finite temperature (in progress)
- ➔ phase transition
- ➔ expanding geometries
- ➔ fermionic d.o.f.
- ...



[Special thanks to LUNA for color pencils]